arXiv:1301.7279v1 [math.PR] 30 Jan 2013

Non-percolation of an asymmetric lilypond model

Christian Hirsch *

January 31, 2013

Abstract

We consider percolation properties of a lilypond model defined by an asymmetric growth-stopping protocol closely related to a model proposed by Daley and Last. More precisely, we show how a sprinkling-based approach can be used to establish non-percolation in a Poisson scenario.

1 Introduction

The classical lilypond model describes a hard-sphere system defined by the following growth-stopping protocol. Start with a point process $X \subset \mathbb{R}^d$ whose atoms serve as germs of a growth process. At time 0 and with the same speed at each element $X_i \in X$ a spherical grain starts to grow. As soon as one such grain touches another grain both cease to grow. This model and various generalizations have been intensively studied for almost 20 years, so that today an entire family of results concerning existence, uniqueness, stabilization and non-percolation is known. We refer the reader to the original articles [2, 3, 4, 5, 6, 7] for details. The purpose of this note is to further advance the completion of this picture by adding a non-percolation result in a lilypond model based on an asymmetric growth-stopping protocol. To be more precise, we consider a model where from each atom of a planar point process a line segment starts to grow in one of the directions $\pm e_1, \pm e_2$ and as soon as a line segment touches an already existing one, the former ceases to grow. This model is an anisotropic variant of the so-called *modified line segment model* discussed in [4].

When using spherical grains the notion of so-called doublets plays a central role, where a *doublet* consists of a pair of ball-shaped grains $B_1, B_2 \subset \mathbb{R}^d$ such that B_1 stops the growth of B_2 and B_2 stops the growth of B_1. This notion allows to subdivide the non-percolation proof provided in [2] into two steps (see also [5] for the earliest non-percolation result). In the first step it is shown that a.s. each connected component of the lilypond model contains at most one doublet. In the second step the a.s. absence of descending chains for homogeneous Poisson point processes is used to show that every connected component also contains at least one doublet. In particular, by mapping each connected component to the midpoint of the two doublet centers we are able construct a locally finite set from the set of connected components in a translation-equivariant way. Therefore an application of the mass-transport principle implies the non-existence of infinite connected components.

In our setting the notion of doublets is replaced by the notion of so-called cycles, where a *cycle* consists of a sequence of line segments $L_1, L_2, \ldots, L_k \subset \mathbb{R}^2$ such that each L_{i+1} $(i = 1, \ldots, k-1)$ stops the growth of L_i and L_1 stops the growth of L_k. As in the setting of spherical grains it is easy to see that any connected component contains at most one cycle. Furthermore, another application of the mass-transport principle proves the absence of infinite connected components containing a cycle. On the other hand, to show that every connected component contains at least one cycle we use the *sprinkling* technique developed in [1]. In other words, we first express the homogeneous Poisson point process $X \subset \mathbb{R}^2$ as

---

*Institute of Stochastics, Ulm University, 89069 Ulm, Germany; E-mail: `christian.hirsch@uni-ulm.de`.

superposition of two independent homogeneous Poisson point processes $X = X^{(1)} \cup X^{(2)}$, where the intensity of $X^{(1)}$ is only slightly smaller than the intensity of X. When considering the lilypond model on $X^{(1)}$, this graph could contain connected components without a cycle, a priori. The idea of the proof is to show that sprinkling the remaining centers $X^{(2)}$ has the effect of stopping every semi-infinite directed path in the lilypond model based on the point process $X^{(1)}$ and that if the sprinkling intensity is chosen sufficiently small, then no additional semi-infinite paths appear. One key step in the formalization of this idea is to combine a stabilization result for the lilypond model at hand with a standard result on dependent percolation [8] to ensure that, except for small exceptional islands in the plane, one has good control on the effects of the sprinkling.

2 Model definition and statement of main result

In this section we fix some notation and provide a formal definition of the asymmetric lilypond model which shall be the topic of our considerations. For ease of presentation we restrict ourselves to an anisotropic model with four fixed directions, although we believe that the general approach presented in this article could be adapted to the isotropic case, as well. For a Borel subset $B \subset \mathbb{R}^2$ we write $\partial B \subset \mathbb{R}^2$ for the topological boundary of B. Furthermore, for $x \in \mathbb{R}^2$ and $s > 0$ we denote by $Q_s(x) = x + [-s/2, s/2]^2$ the s-square centered at x and by $Q_s^m(x) = Q_s(x) \times M$ the product of this square with the mark space $M = \{\pm e_1, \pm e_2\}$ which is to be identified with the set of possible directions of growth. Similarly, we write $\mathbb{R}^{2,m} = \mathbb{R}^2 \times M$ for the product of $\mathbb{R}^2$ with the mark space M. A locally finite subset $\varphi \subset \mathbb{R}^{2,m}$ will usually be interpreted as a pair consisting of a locally finite set $\varphi \subset \mathbb{R}^2$ and a function $dir : \varphi \to M$ indicating the direction of growth. Furthermore, for $s > 0$, $z \in \mathbb{Z}^2$ and $\varphi \subset \mathbb{R}^{2,m}$ locally finite we denote by $\varphi_{z,s} = \varphi \cap Q_s^m(sz)$ the restriction of φ to the (marked) square $Q_s^m(sz)$. If the value of s is understood (which will usually be the case) we also simply write φ_z instead of $\varphi_{z,s}$. Having fixed some notation we now provide a precise definition of the lilypond model.

Definition 2.1. Let $\varphi \subset \mathbb{R}^{2,m}$ be locally finite. For a given function $f : \varphi \to [0, \infty]$ define the associated *grain-endpoint function* $g : \varphi \to \mathbb{R}^2 \cup \{\infty\}$ and *grain function* $I : \varphi \to \mathcal{P}\left(\mathbb{R}^2\right)$ by

$$g(x) = \begin{cases} x + f(x)dir(x) & \text{if } f(x) \neq \infty, \\ \infty & \text{if } f(x) = \infty, \end{cases}$$

and

$$I(x) = \begin{cases} [x, g(x)) & \text{if } f(x) \neq \infty \\ x + [0, \infty)dir(x), & \text{if } f(x) = \infty, \end{cases}$$

respectively. We say that f constitutes a *lilypond radius* if the following two items are satisfied.

(i) *hard-core property:* $I(x) \cap I(y) = \emptyset$ for all $x, y \in \varphi$ with $x \neq y$.

(ii) *existence of stopping neighbors:* for all $x \in \varphi$ with $f(x) \neq \infty$ there exists a unique $y \in \varphi$ with $g(x) \in I(y)$ and, additionally, the inequality $|g(x) - y| < f(x)$ holds.

If there exists a unique such function f, we denote it by f_φ (and the associated g and I by g_φ and I_φ). Furthermore, we then also define a function $h_\varphi : \varphi \to \varphi$ mapping x to its unique stopping neighbor.

A general principle which has proven to be applicable in a large number of lilypond-type models states that existence and uniqueness of the lilypond radius hold if the underlying locally finite set does not admit a suitable form of descending chain. This is also true for the lilypond model at hand. To be more precise, we denote by $\mathbb{N}'$ the family of all locally finite subsets $\varphi \subset \mathbb{R}^{2,m}$ such that

(i) φ does not admit an infinite anisotropic descending chain in the sense of Definition 4.1, and

(ii) $(x-y)/|x-y|_2 \notin \{\pm e_1, \pm e_2, (\pm e_1 \pm e_2)/\sqrt{2}\}$ for all $x, y \in \varphi$ with $x \neq y$.

Then using the same arguments as e.g. in [4] one can prove that for all $\varphi \in \mathbb{N}'$ there exists a unique lilypond radius $f_\varphi : \varphi \to [0,\infty]$. Furthermore, if $X \subset \mathbb{R}^{2,m}$ denotes an independently and uniformly marked homogeneous Poisson point process, then $X \in \mathbb{N}'$ and $f_X(X) \subset [0,\infty)$ a.s.

Remark 1. Let $\varphi^{(1)} \subset \varphi^{(2)} \subset \mathbb{R}^{2,m}$ be locally finite with $\varphi^{(1)}, \varphi^{(2)} \in \mathbb{N}'$. Then for any $x \in \varphi^{(1)}$ the relation $f_{\varphi^{(1)}}(x) = f_{\varphi^{(2)}}(x)$ is equivalent to $h_{\varphi^{(1)}}(x) = h_{\varphi^{(2)}}(x)$.

For $i \geq 0$, $\varphi \in \mathbb{N}'$ and $x \in \varphi$, the *i-th lilypond descendant of x* is defined recursively by $h^i_\varphi(x) = x$ if $i = 0$ and $h^i_\varphi(x) = h_\varphi\left(h^{i-1}_\varphi(x)\right)$ if $i \geq 1$. We also denote by $h^\infty_\varphi(x) = \left\{h^i_\varphi(x) \mid i \geq 0\right\}$ the set of all lilypond descendants of x. The main theorem of this article concerns the a.s. finiteness of $h^\infty_X(x)$ in the Poisson case.

Theorem 2.2. *Let $X \subset \mathbb{R}^{2,m}$ denotean independently and uniformly M-marked homogeneous Poisson point process with intensity 1. Then with probability 1 the set $h^\infty_X(x)$ is finite for all $x \in X$.*

In the following we frequently consider the behavior of the lilypond model inside squares of the form $Q_s(sz)$ for some $s > 0$ and $z \in \mathbb{Z}^d$. Therefore, it is convenient to introduce some notation for germs $x \in \varphi$ whose associated grain $I_\varphi(x)$ enters or leaves $Q_s(sz)$. More precisely, we denote by $\partial^{in}_\varphi Q_s(sz) = \{x \in \varphi \setminus Q^m_s(sz) : I_\varphi(x) \cap Q_s(sz) \neq \emptyset\}$ the set of all $x \in \varphi \setminus Q^m_s(sz)$ entering $Q_s(sz)$ and by $\partial^{out}_\varphi Q_s(sz) = \{x \in \varphi_{z,s} : I_\varphi(x) \not\subset Q_s(sz)\}$ the set of all $x \in \varphi_{z,s}$ leaving $Q_s(sz)$.

3 Proof of Theorem 2.2

In this section we discuss the main ideas for the proof of Theorem 2.2. Our goal is to present the general framework while proofs of important technical lemmas are postponed to Section 4. In the following we always assume that $X \subset \mathbb{R}^{2,m}$ denotes a homogeneous Poisson point process on $\mathbb{R}^2$ with intensity 1 and which is independently marked with uniform marks in M.

As indicated in Section 1 the proof of Theorem 2.2 is based on a sprinkling argument. The sprinkling has to be analyzed in a rather delicate way because two essential properties must be satisfied. On the one hand, we have to guarantee that if $x \in X^{(1)}$ is an element with $\# h^\infty_{X^{(1)}}(x) = \infty$, then after the sprinkling we must have $\# h^\infty_X(x) < \infty$. On the other the sprinkling should not influence the lilypond model too strongly, since for all $x \in X^{(1)}$ with $\# h^\infty_{X^{(1)}}(x) < \infty$ it has to be ensured that also after the sprinkling $\# h^\infty_X(x)$ is still finite. We solve this problem by adding the sprinkling in two steps. First, we construct a point process $X^{(3)} \subset \mathbb{R}^2$ with $X^{(1)} \subset X^{(3)} \subset X$ by adding only those germs of $X^{(2)}$ for which we have little knowledge of as how their addition would affect the existing lilypond model. In a second step we add the remaining germs of $X^{(2)}$ for which we have precise information about their impact on the already existing model. To implement this strategy, an external stabilization result will constitute a crucial intermediate step.

First we decompose the homogeneous Poisson point process $X \subset \mathbb{R}^{2,m}$ as superposition $X = X^{(1)} \cup X^{(2)}$ of two independently and uniformly M-marked homogeneous Poisson point processes $X^{(1)}$ and $X^{(2)}$, which are independent and have intensities $1-\lambda, \lambda > 0$, respectively, where typically $\lambda \ll 1$ (in the following we assume $\lambda \in (0, 1/4)$; the precise value of λ is fixed in Lemma 3.7). Next we discretize $\mathbb{R}^2$ and define a site-percolation process consisting of those discretized squares with four nice properties that turn out to be useful when trying to control the effect of the sprinkling.

Definition 3.1. Let $s \geq 1$ (where typically $s \gg 1$) and put $\delta = s^{-4}$. We say that $z \in \mathbb{Z}^2$ is *s-good* iff it satisfies the following four properties.

(i) *external stabilization:* $\partial^{in}_{X^{(1)} \cap Q^m_{3s}(sz)} Q_s(sz) = \partial^{in}_{X^{(1)} \cap Q^m_{3s}(sz) \cup \xi} Q_s(sz)$ for all finite $\xi \subset \mathbb{R}^{2,m} \setminus Q^m_{3s}(sz)$.

(ii) *δ-separatedness:* $X_z^{(1)} \cup \partial^{in}_{X^{(1)}} Q_s(sz) \cup \{sz + \frac{s}{2}M\}$ is δ-separated, i.e., $|\pi_k(x) - \pi_k(y)| \geq \delta$ for all $x, y \in X_z^{(1)} \cup \partial^{in}_{X^{(1)}} Q_s(sz) \cup \{sz + \frac{s}{2}M\}$ with $x \neq y$ and all $k \in \{1,2\}$, where $\pi_k : \mathbb{R}^{2,m} \to \mathbb{R}$ denotes the projection to the k-th spatial coordinate.

(iii) *absence of trespassing segments:* $g_{X^{(1)}}(x) \in Q_s(sz)$ for all $x \in \partial^{in}_{X^{(1)}} Q_s(sz)$.

(iv) *existence of interior segments:* $X_z^{(1)} \neq \partial^{out}_{X^{(1)}} Q_s(sz)$.

Remark 2. It is easy to check that for all locally finite $\xi \subset \mathbb{R}^{2,m} \setminus Q^m_{3s}(sz)$ such that $\xi \cup X^{(1)} \in \mathbb{N}'$ there exists a *finite* $\xi' \subset \mathbb{R}^{2,m} \setminus Q^m_{3s}(sz)$ such that $\partial^{in}_{X^{(1)} \cap Q^m_{3s}(sz) \cup \xi'} Q_s(sz) = \partial^{in}_{X^{(1)} \cap Q^m_{3s}(sz) \cup \xi} Q_s(sz)$

As we will see later (in Lemma 4.10) if item (i) is satisfied, then the validity of items (ii), (iii), (iv) only depends on $X \cap Q^m_{3s}(sz)$. In particular, the site process of s-good sites is 3-dependent. If we can show that, by increasing s, the probability that o is s-good can be made as close to 1 as desired, then [8, Theorem 0.0] is applicable. In particular, the set of s-bad sites can be dominated from above by a subcritical Bernoulli site-percolation process. This assertion is the content of the following result whose proof is postponed to Section 4.2.

Proposition 3.2. $\lim_{s\to\infty} \mathbb{P}(o \textit{ is } s\textit{-good}) = 1$.

Recall that we want to use the sprinkled germs $X^{(2)}$ to ensure that $\# h^\infty_X(x) < \infty$ even for those elements $x \in X^{(1)}$ for which $\# h^\infty_{X^{(1)}}(x) = \infty$ cannot be excluded a priori. In other words the sprinkled germs should be capable of stopping all semi-infinite paths in the lilypond model on $X^{(1)}$. Next we introduce special local configurations in $X^{(2)}$ which play an important role in this stopping mechanism.

Definition 3.3. Let $\delta > 0$ and $x \in \mathbb{R}^2$. We say that $D = \{x_1, \dots, x_4\} \subset Q^m_\delta(x)$ forms an *(x, δ)-cycle* if the following items are satisfied.

(i) $dir(x_j) = \rho_{\pi/2} dir(x_{j+1})$ for all $j \in \{1, \dots, 4\}$ (with indices j considered cyclically *mod* 4), where $\rho_{\pi/2} : \mathbb{R}^2 \to \mathbb{R}^2$ denotes rotation by $\pi/2$,

(ii) $g_D(D) \subset Q_\delta(x)$,

(iii) $g_D(x_j) \in I_D(x_{j+1})$ for all $j \in \{1, \dots, 4\}$ and

(iv) $\min(\langle x - x_j, dir(x_j)\rangle, \langle x - x_j, dir(x_{j+1})\rangle) > 0$ for all $j \in \{1, \dots, 4\}$

If D forms a (x', δ)-cycle for some $x' \in \mathbb{R}^2$, we say that D forms a δ-cycle. Finally, write $A^{(1)}_{x,\delta}(\xi)$ for the event that the configuration of $\xi \cap Q_\delta(x)$ consists precisely of one (x, δ)-cycle. See Figure 1 for an illustration of an (x, δ)-cycle.



Figure 1: Example of an (x, δ)-cycle

An important feature of δ-cycles is their strong external stability property which is established in the following lemma.

Lemma 3.4. *Let $\delta > 0$, $x \in \mathbb{R}^2$ and $\varphi \in \mathbb{N}'$ be such that $Q_{3\delta}(x) \cap \varphi = \emptyset$. Furthermore, let $D = \{x_1, \ldots, x_4\} \subset \mathbb{R}^{2,m}$ form an (x, δ)-cycle. Then $f_{D\cup\varphi}(x_i) = f_D(x_i)$ for all $i \in \{1, \ldots, 4\}$.*

Proof. Suppose there exists $i \in \{1, \ldots, 4\}$ with $f_{D\cup\varphi}(x_i) \neq f_D(x_i)$. By the hard-core property we see that we cannot have $f_{D\cup\varphi}(x_j) \geq f_D(x_j)$ for all $j \in \{1, \ldots, 4\}$ and we choose $j_1 \in \{1, \ldots, 4\}$ with $f_{D\cup\varphi}(x_{j_1}) < f_D(x_{j_1})$. Note that

$$\begin{aligned}|h_{D\cup\varphi}(x_{j_1}) - x|_\infty &\leq |h_{D\cup\varphi}(x_{j_1}) - x_{j_1}|_\infty + |x_{j_1} - x|_\infty \\ &\leq f_{D\cup\varphi}(x_{j_1}) + \delta/2 \\ &\leq 3\delta/2,\end{aligned}$$

so that $h_{D\cup\varphi}(x_{j_1}) \in Q_{3\delta}(x)$, violating the assumption $Q_{3\delta}(x) \cap \varphi = \emptyset$. □

The notion of (x, δ)-cycles can be used to create configurations of $X_z^{(2)}$ satisfying the relation $\# h^\infty_{X^{(1)} \cup X_z^{(2)}}(x) < \infty$ for all $x \in \partial^{in}_{X^{(1)}} Q_s(sz)$. To make this more precise, we consider the following definition.

Definition 3.5. Let $s > 0$, $z \in \mathbb{Z}^2$, $\xi \subset Q_s^m(sz)$ finite and $\varphi \in \mathbb{N}'$. Then we put $\delta = s^{-4}$, $M_1 = \left\{x + \frac{3\delta}{8} dir(x) : x \in \varphi_z \setminus \partial^{out}_\varphi Q_s(sz)\right\}$ and $M_2 = \left\{P_x + \frac{3\delta}{8} dir(x) : x \in \partial^{in}_\varphi Q_s(sz)\right\}$, where P_x denotes the first intersection point of $I_\varphi(x)$ and $\partial Q_s(sz)$. Then we define $A^{(2)}_{s,z}(\varphi, \xi)$ to be the intersection of the events $\left\{\xi \subset (M_1 \cup M_2) \oplus Q_{\delta/16}(o)\right\}$ and $\bigcap_{x \in M_1 \cup M_2} A^{(1)}_{x,\delta/16}(\xi)$. If the value of s is understood, we also write $A^{(2)}_z(\varphi, \xi)$ instead of $A^{(2)}_{s,z}(\varphi, \xi)$. See Figure 2 for an illustration of the effect on the lilypond model when adding a set of germs ξ satisfying $A^{(2)}_z(\varphi, \xi)$.


(a) Configuration before addition of ξ (b) Configuration after addition of ξ

Figure 2: Impact of the addition of ξ with $A^{(2)}(\varphi, \xi)$

Informally speaking, configurations $\xi \subset Q_s^m(sz)$ satisfying $A^{(2)}_z(\varphi, \xi)$ have the effect of stopping every grain entering $Q_s(sz)$ while ensuring that $\partial^{out}_\varphi Q_s(sz)$ is left unchanged, so that addition of the sprinkling does not have any effects on the configuration of the lilypond model outside of $Q_s(sz)$. More precise results are stated and proven in Section 4.3.

Next we introduce a discrete site process which allows us to determine whether there is either

(i) no sprinkling inside the corresponding square, or

(ii) a sprinkling as described above, or

(iii) some other sprinkling.

To be more precise, define a $\{0,1,2\}$-valued site process $\{Y_z\}_{z\in\mathbb{Z}^2}$ as follows. If $X_z^{(2)} = \emptyset$ and z is s-good, then we put $Y_z = 0$. Next we put $Y_z = 1$ if z is s-good and the event $A_z^{(2)}\left(X^{(1)}, X_z^{(2)}\right)$ occurs. If neither $Y_z = 0$ nor $Y_z = 1$, then we put $Y_z = 2$. Note that conditioned on $X^{(1)}$ the site process $\{Y_z\}_{z\in\mathbb{Z}^2}$ forms an inhomogeneous Bernoulli-site process.

Next we observe that given $X^{(1)}$ the probability of seeing a useful sprinkling in a given square is bounded from below by a positive constant, as long as the respective square is s-good.

Lemma 3.6. *There exists $p > 0$ such that $\mathbb{P}\left(Y_z = 1 \mid X^{(1)}\right) \ge p 1_{z \text{ is } s\text{-good}}$ for all $z \in \mathbb{Z}^2$. In other words, p may depend on s and λ, but* not *on z or a given realization of $X^{(1)}$.*

Proof. Let z be s-good and let $M_1, M_2 \subset Q_s(sz)$ be as in the definition of the event $A_z^{(2)}\left(X^{(1)}, X_z^{(2)}\right)$. We conclude from δ-separatedness that for all $x_1, x_2 \in M_1 \cup M_2$ with $x_1 \neq x_2$ we have $Q_{\delta/16}(x_1) \cap Q_{\delta/16}(x_2) = \emptyset$. In particular, for any $x \in M_1 \cup M_2$ the event $A^{(1)}_{x,\delta/16}\left(X_z^{(2)}\right)$ is independent of the family of events $A^{(1)}_{y,\delta/16}\left(X_z^{(2)}\right)$ for $y \in M_1 \cup M_2$ with $y \neq x$. Furthermore, from δ-separatedness we also conclude $|M_1| \le s^2\delta^{-2}$ and $|M_2| \le 4s\delta^{-1}$. Finally, note that $\mathbb{P}\left(X_z^{(2)} \subset (M_1 \cup M_2) \oplus Q_{\delta/16}(o)\right) \ge \mathbb{P}\left(X_z^{(2)} \cap Q_s(sz) = \emptyset\right)$. Hence, we may choose $p = \mathbb{P}\left(X_z^{(2)} \cap Q_s(o) = \emptyset\right)\mathbb{P}\left(A^{(1)}_{o,\delta/16}\left(X_z^{(2)}\right)\right)^{s^2\delta^{-2}+4s\delta^{-1}}$. □

In the next step we identify a large set of sites for which we have good control over the effect of the sprinkling. We recursively construct sets of revealed sites $\left(R^{(i)}\right)_{i\ge 0}$, $R^{(i)} \subset \mathbb{Z}^2$ and bad sites $\left(B^{(i)}\right)_{i\ge 0}$, $B^{(i)} \subset \mathbb{Z}^2$ as follows. Initially, put $R^{(0)} = B^{(0)} = \left\{z \in \mathbb{Z}^2 : Y_z = 2\right\}$. Now suppose $i \ge 0$ and that $R^{(i)} \subset \mathbb{Z}^2$ as well as $B^{(i)} \subset \mathbb{Z}^2$ have already been constructed. For $z \in \mathbb{Z}^2$ write $S(z) = \{z' \in \mathbb{Z}^2 : |z - z'|_\infty \le 2\}$. Choose the closest bad site $z \in \mathbb{Z}^2$ to the origin with the property that its neighborhood $S(z)$ is not already completely revealed, i.e., $z \in B^{(i)}$ but $S(z) \not\subset R^{(i)}$ (if several sites have this property, we choose the lexicographically smallest one). Put $R^{(i+1)} = R^{(i)} \cup S(z)$ and $B^{(i+1)} = B^{(i)} \cup \{z' \in S(z) : X_{z'}^{(2)} \neq \emptyset\}$. Finally, put $R = \bigcup_{i\ge 0} R^{(i)}$ and see Figure 3 for an illustration of the construction of the set R.


(a) Initial set of bad sites (b) New bad sites revealed (c) No new bad sites revealed

Figure 3: Construction of R

We first note that for $\lambda \ll 1$ only very few sites are revealed at all.

Lemma 3.7. *There exist $s > 1$ and $\lambda \in (0, 1/4)$ such that with probability 1 the set $R \subset \mathbb{Z}^2$ is dominated from above by a sub-critical Bernoulli site-percolation process.*

Proof. We first note that $R \subset Y^{(a)} \cup Y^{(b)}$, where $Y^{(a)} \subset \mathbb{Z}^2$ denotes the set of sites $z \in \mathbb{Z}^2$ whose neighborhood $S(z)$ contains a site which is not s-good and where $Y^{(b)} \subset \mathbb{Z}^2$ consists of those $z \in \mathbb{Z}^2$ with $X^{(2)} \cap Q_{5s}(sz) \neq \emptyset$. We note that $Y^{(a)}$ and $Y^{(b)}$ are 7-dependent site processes which are independent of each other, so that the claim follows from Proposition 3.2 and [8, Theorem 0.0]. □

In the remaining part of this section we fix $s > 1$ and $\lambda \in (0, 1/4)$ so that the conclusion of Lemma 3.7 holds. Then we define $X^{(3)} = X^{(1)} \cup \big(X^{(2)} \cap (Q_s(o) \oplus R)\big)$. In other words, to create the original Poisson point process X from $X^{(3)}$ we only have to add the sprinkling $X^{(2)}$ in the unrevealed region $Q_s(o) \oplus R^c$, where $R^c = \mathbb{Z}^2 \setminus R$ denotes the complement of R in $\mathbb{Z}^2$. We first note an auxiliary result which allows us to compare the lilypond model on $X^{(1)}$ to the lilypond model on $X^{(3)}$ and whose proof is postponed to Section 4.3.

Lemma 3.8. *For all $z \in R^c$ the following identities hold.*

(i) $\partial^{in}_{X^{(3)}} Q_s(sz) = \partial^{in}_{X^{(1)}} Q_s(sz)$.

(ii) $f_{X^{(3)}}(x) = f_{X^{(1)}}(x)$ *for all* $x \in X^{(3)}$ *with* $\{g_{X^{(3)}}(x), g_{X^{(1)}}(x)\} \cap Q_s(sz) \neq \emptyset$.

In order to compare $h^{\infty}_{X^{(3)}}$ and h^{∞}_X it is convenient to understand the effect of adding the sprinkled nodes $X^{(2)} \setminus (Q_s(o) \oplus R)$ in a step-by-step manner. However, as the proof of the following result is rather technical, it is also deferred to Section 4.3 .

Lemma 3.9. *Let $\{z_1, z_2, \ldots\}$ be an arbitrary enumeration of R^c and put $X^{(2,j)} = \bigcup_{i=1}^{j} X^{(2)}_{z_i}$ as well as $X^{(3,j)} = X^{(3)} \cup X^{(2,j)}$. Then for all $j \geq 0$*

(i) $\partial^{in}_{X^{(3,j)}} Q_s(sz) = \partial^{in}_{X^{(1)}} Q_s(sz)$ *for all* $z \in R^c$,

(ii) $f_{X^{(3,j)}}(x) = f_{X^{(3)}}(x)$ *for all* $z \in R$ *and all* $x \in X^{(3)}$ *with* $g_{X^{(3)}}(x) \in Q_s(sz)$,

(iii) $f_{X^{(3,j)}}(x) = f_{X^{(2,j)}_z}(x)$ *for all* $z \in R^c$ *and all* $x \in X^{(2,j)}_z$

(iv) $f_{X^{(3,j)}}(x) = f_{X^{(1)}}(x)$ *for all* $z \in R^c$ *with* $X^{(2,j)}_z = \emptyset$ *and all* $x \in X^{(3)}$ *with* $g_{X^{(3)}}(x) \in Q_s(sz)$ *and*

(v) $h_{X^{(3,j)}}(x) \in X^{(2)}_z$ *for all* $z \in R^c$ *with* $X^{(2,j)}_z \neq \emptyset$ *and all* $x \in X^{(3)}$ *with* $g_{X^{(3)}}(x) \in Q_s(sz)$.

The following deterministic result allows us to pass to the limit $j \to \infty$.

Lemma 3.10. *Let $\varphi^{(0)} \subset \varphi^{(1)} \subset \cdots \subset \varphi = \bigcup_{i \geq 0} \varphi^{(i)} \subset \mathbb{R}^{2,m}$ be such that $\varphi \in \mathbb{N}'$ and suppose that for all $i \geq 0$ there exists $n(i) \geq i$ with $f_{\varphi^{(n(i))}}(x) = f_{\varphi^{(j)}}(x)$ for all $j \geq n(i)$ and all $x \in \varphi^{(i)}$. Then $f_{\varphi^{(n(i))}}(x) = f_{\varphi}(x)$ for all $i \geq 0$ and all $x \in \varphi^{(i)}$.*

Proof. It is easy to check that the function $f : \varphi \to [0, \infty]$, $x \mapsto f_{\varphi^{(n(i))}}(x)$ if $x \in \varphi^{(i)}$ is both well-defined and a lilypond radius. ☐

Using Lemma 3.10 we therefore obtain the following corollary.

Corollary 3.11. *For all $j \geq 0$ we have*

(i) $\partial^{in}_{X} Q_s(sz) = \partial^{in}_{X^{(1)}} Q_s(sz)$ *for all* $z \in R^c$,

(ii) $f_X(x) = f_{X^{(3)}}(x)$ *for all* $z \in R$ *and all* $x \in X^{(3)}$ *with* $g_{X^{(3)}}(x) \in Q_s(sz)$,

(iii) $f_X(x) = f_{X^{(2)}_z}(x)$ *for all* $z \in R^c$ *and all* $x \in X^{(2)}_z$,

(iv) $f_X(x) = f_{X^{(1)}}(x)$ *for all* $z \in R^c$ *with* $X^{(2)}_z = \emptyset$ *and all* $x \in X^{(3)}$ *with* $g_{X^{(3)}}(x) \in Q_s(sz)$,

(v) $h_X(x) \in X^{(2)}_z$ *for all* $z \in R^c$ *with* $X^{(2)}_z \neq \emptyset$ *and all* $x \in X^{(3)}$ *with* $g_{X^{(3)}}(x) \in Q_s(sz)$.

After this preliminary work the following comparison result between $h^{\infty}_{X^{(3)}}$ and h^{∞}_X is established straightforwardly.

Lemma 3.12. *If $x \in X$ is such that $\# h^{\infty}_X(x) = \infty$, then $h^{\infty}_X(x) \subset X^{(3)}$ and $h^i_X(x) = h^i_{X^{(3)}}(x)$ for all sufficiently large $i \geq 0$.*

Proof. Before providing a formal proof of this statement we consider a heuristic explanation. Let $x \in X$ be such that $\# h^{\infty}_X(x) = \infty$. Observe that $X^{(3)}$ and X only differ in squares $Q_s(sz)$ with $z \in R^c$ and $Y_z = 1$. In particular, if the polygonal path $\gamma = \bigcup_{i \geq 0} I_X\left(h^i_X(x)\right) \subset \mathbb{R}^2$ induced by $h^{\infty}_X(x)$ does not intersect such squares we have $x \in X^{(3)}$ and $h^i_X(x) = h^i_{X^{(3)}}(x)$ for all $i \geq 0$. On the other hand if γ enters such a square $Q_s(sz)$ then we conclude from the occurrence of $A^{(2)}\left(X^{(3)}, X_z^{(2)}\right)$ that γ cannot be semi-infinite but must end inside this square.

To make this formal, consider $x \in X$ with $\# h_X^\infty(x) = \infty$. If there exists $i \geq 0$ with $h_X^i(x) \in X_z^{(2)}$ for some $z \in R^c$, then item (iii) of Corollary 3.11 shows $h_X^{i+5}(x) = h_X^{i+1}(x)$, thereby contradicting $\# h_X^\infty(x) = \infty$. Hence, $h_X^\infty(x) \subset X^{(3)}$. Next, we observe that there exists $i_0 \geq 1$ such that for all $i \geq i_0$ if $z \in \mathbb{Z}^2$ is chosen such that $g_X(h_X^i(x)) \in Q_s(sz)$, then we have $z \in R$ or $Y_z \neq 1$. Indeed, otherwise choose any $i \geq 0$ with $h_X^i(x) \in \partial_X^{in} Q_s(sz)$ and $g_X(h_X^i(x)) \in Q_s(sz)$ for some $z \in R^c$ satisfying $Y_z = 1$. By item (i) of Corollary 3.11 we obtain $h_X^i(x) \in \partial_{X^{(1)}}^{in} Q_s(sz)$, so that Lemma 3.8 and absence of trespassing segments imply $g_{X^{(3)}}(h_X^i(x)) = g_{X^{(1)}}(h_X^i(x)) \in Q_s(sz)$. By item (v) of Corollary 3.11 we conclude $h_X(h_X^i(x)) \in X_z^{(2)}$ contradicting $h_X^\infty(x) \subset X^{(3)}$. This completes the proof of the existence of i_0 as described above.

Finally, we show $f_X\left(h_X^i(x)\right) = f_{X^{(3)}}\left(h_X^i(x)\right)$ for all $i \geq i_0$ (which implies $h_X\left(h_X^i(x)\right) = h_{X^{(3)}}\left(h_X^i(x)\right)$ for all $i \geq i_0$ by Remark 1). We prove this assertion by distinguishing two cases. If $g_{X^{(3)}}(h_X^i(x)) \in Q_s(sz)$ with $z \in R$, then the assertion follows immediately from item (ii) of Corollary 3.11. Otherwise item (iv) of the same corollary and Lemma 3.8 imply $f_X(h_X^i(x)) = f_{X^{(1)}}(h_X^i(x)) = f_{X^{(3)}}(h_X^i(x))$. ☐

Proof of Theorem 2.2. Assume the contrary for the sake of deriving a contradiction. Then by Lemma 3.12 there exists $x \in X^{(3)}$ such that $\# h_{X^{(3)}}^\infty(x) = \infty$ and $h_X(h_{X^{(3)}}^i(x)) \in X^{(3)}$ for all $i \geq 0$. We denote this event by $A_x^{(3)}$. Then it suffices to verify $\mathbb{P}\left(A_x^{(3)} \mid X^{(1)}, X^{(3)}\right) = 0$ for all $x \in X^{(3)}$.

If $\# h_{X^{(3)}}^\infty(x) = \infty$ then we note that by Lemma 3.7 there exist infinitely many $z_1, z_2, \ldots \in R^c$ and $i_1, i_2, \ldots \geq 1$ with $h_{X^{(3)}}^{i_j}(x) \in \partial_{X^{(3)}}^{in} Q_s(sz_j)$ for all $j \geq 1$. Furthermore, observe that conditioned on $X^{(1)}$ and $X^{(3)}$ the restriction of the site process $\{Y_z\}_{z \in \mathbb{Z}^2}$ to R^c defines an inhomogeneous $\{0,1\}$-valued Bernoulli site process such that for $z \in R^c$ the (conditional) probability of the event $\{Y_z = 1\}$ is given by $\mathbb{P}\left(Y_z = 1 \mid Y_z \in \{0,1\}, X^{(1)}\right)$. In particular the events $\left\{Y_{z_j} = 1\right\}$ occur independently given $X^{(1)}$ and $X^{(3)}$ and by Lemma 3.6 we have $\mathbb{P}\left(Y_z = 1 \mid Y_z \in \{0,1\}, X^{(1)}\right) \geq p > 0$ a.s. Therefore, with probability 1 there exists $j_0 \geq 1$ with $Y_{z_{j_0}} = 1$. In particular, there exists a δ-cycle $D = \left\{X_{i_j}^{(2)}\right\}_{1 \leq j \leq 4}$ with $h_X\left(h_{X^{(3)}}^{i_{j_0}}(x)\right) \in D$ and $D \cap X^{(3)} = \emptyset$. This shows $\mathbb{P}\left(A_x^{(3)} \mid X^{(1)}, X^{(3)}\right) = 0$, thereby completing the proof of Theorem 2.2. ☐

4 Technical lemmas

4.1 Preliminaries on anisotropic descending chains

As in the classical setting discussed in [2] the notion of descending chains – or rather a variant thereof – is closely related to a variety of properties of the considered lilypond model.

Definition 4.1. Let $b > 0$ and let $\varphi \subset \mathbb{R}^2$ be locally finite. We say that a (finite or infinite) set $\{x_i\}_{i \geq 1} \subset \varphi$ forms a *b-bounded anisotropic descending chain* if $|x_1 - x_2|_\infty \leq b$ and $|x_i - x_{i+1}|_\infty < |x_{i-1} - x_i|_\infty$ for all $i \geq 2$. We simply say $\{x_i\}_{i \geq 1} \subset \varphi$ forms an *anisotropic descending chain* if it forms a *b-bounded anisotropic descending chain* for some $b > 0$.

Lemma 4.2. *Let $\varphi^{(1)}, \varphi^{(2)} \in \mathbb{N}'$ and suppose $x_1 \in \varphi^{(1)} \cap \varphi^{(2)}$ is such that $f_{\varphi^{(1)}}(x_1) < f_{\varphi^{(2)}}(x_1)$. Furthermore, define recursively $x_{i+1} \in \varphi^{(1)} \cup \varphi^{(2)}$ by*

$$x_{i+1} = \begin{cases} h_{\varphi^{(1)}}(x_i) & \text{if } x_i \in \varphi^{(1)} \\ x_i & \text{else} \end{cases}$$

if i is odd and by

$$x_{i+1} = \begin{cases} h_{\varphi^{(2)}}(x_i) & \text{if } x_i \in \varphi^{(2)} \\ x_i & \text{else} \end{cases}$$

if i is even. Finally, put $i_0 = \min\{i \geq 1 : x_i \notin \varphi^{(1)} \cap \varphi^{(2)}\}$. Then $(x_i)_{1 \leq i \leq i_0}$ constitutes an $f_{\varphi^{(1)}}(x_1)$-bounded anisotropic descending chain of pairwise distinct elements. In particular, the absence of infinite anisotropic descending chains in $\varphi^{(1)}$ and $\varphi^{(2)}$ implies $i_0 < \infty$. Additionally $f_{\varphi^{(1)}}(x_i) < f_{\varphi^{(2)}}(x_i)$ if $i \in \{1, \dots i_0 - 1\}$ is odd and $f_{\varphi^{(2)}}(x_i) < f_{\varphi^{(1)}}(x_i)$ if $i \in \{1, \dots, i_0 - 1\}$ is even.

Proof. The relation $|x_1 - x_2|_\infty = f_{\varphi^{(1)}}(x_1)$ follows immediately from the definition of stopping neighbors. Next, assume that $i \geq 2$ is odd. From $x_i = h_{\varphi^{(2)}}(x_{i-1})$ and $x_{i+1} = h_{\varphi^{(1)}}(x_i)$ we conclude $|x_i - x_{i-1}|_\infty = f_{\varphi^{(2)}}(x_{i-1})$ and $|x_{i+1} - x_i|_\infty = f_{\varphi^{(1)}}(x_i)$. Furthermore, by induction we have $f_{\varphi^{(2)}}(x_{i-1}) < f_{\varphi^{(1)}}(x_{i-1})$, so that

$$f_{\varphi^{(1)}}(x_i) < \left|x_i - g_{\varphi^{(2)}}(x_{i-1})\right| < \min\left(f_{\varphi^{(2)}}(x_{i-1}), f_{\varphi^{(2)}}(x_i)\right). \tag{1}$$

Using similar arguments, we see that $|x_{i+1} - x_i|_\infty < |x_i - x_{i-1}|_\infty$ and $f_{\varphi^{(2)}}(x_i) < f_{\varphi^{(1)}}(x_i)$ hold when $i \geq 2$ is even.

It remains to show that the $\{x_i\}_{1 \leq i \leq i_0}$ are pairwise disjoint and by the definition of i_0 it suffices to prove this claim for $\{x_i\}_{1 \leq i < i_0}$. Furthermore, as $f_{\varphi^{(1)}}(x_i) < f_{\varphi^{(2)}}(x_i)$ if $i \in \{1, \dots, i_0 - 1\}$ is odd and $f_{\varphi^{(2)}}(x_i) < f_{\varphi^{(1)}}(x_i)$ if $i \in \{1, \dots, i_0 - 1\}$ is even, it suffices to consider the pairwise disjointness of the points in $\{x_i\}_{\substack{1 \leq i < i_0 \\ i \text{ even}}}$ and the points in $\{x_i\}_{\substack{1 \leq i < i_0 \\ i \text{ odd}}}$ separately. We consider for instance the case of even $i \in \{1, \dots, i_0 - 1\}$. Then we note that an application of (1) and its analog for even parity show

$$f_{\varphi^{(2)}}(x_i) > f_{\varphi^{(1)}}(x_{i+1}) > f_{\varphi^{(2)}}(x_{i+2}),$$

so that by induction we obtain $f_{\varphi^{(2)}}(x_i) > f_{\varphi^{(2)}}(x_j)$ for all $i, j \in \{1, \dots, i_0 - 1\}$ such that i, j are even and $i < j$. □

To obtain bounds for the probability of occurrence of long anisotropic descending chains the following auxiliary computation is useful.

Lemma 4.3. *Let $b, k \geq 0$ and $x_0 \in \mathbb{R}^2$. Then $\int_{\mathbb{R}^2} 1_{|x - x_0|_\infty \leq b} |x - x_0|_\infty^{2k} \mathrm{d}x = \frac{4}{k+1} b^{2k+2}$.*

Proof. Without loss of generality we may assume $x_0 = o$ and we write $x = (\xi_1, \xi_2)$. Then by symmetry

$$\begin{aligned}\int 1_{|x|_\infty \leq b} |x|_\infty^{2k} \mathrm{d}x &= 4 \int_0^b \int_0^b \max\left(\xi_1, \xi_2\right)^{2k} \mathrm{d}\xi_2 \mathrm{d}\xi_1 \\ &= 8 \int_0^b \xi_1^{2k} \int_0^{\xi_1} \mathrm{d}\xi_2 \mathrm{d}\xi_1 \\ &= 8 \int_0^b \xi_1^{2k+1} \mathrm{d}\xi_1 \\ &= \frac{8}{2k+2} b^{2k+2}.\end{aligned}$$

□

Using Lemma 4.3 we obtain the following result.

Lemma 4.4. *Let $n \geq 1$, $b, s > 0$ and let $X \subset \mathbb{R}^2$ be a homogeneous Poisson point process with intensity $\lambda > 0$. Denote by $A^{(4)}$ the event that there exist pairwise disjoint $X_0, \dots, X_n \in X$, with $X_0 \in Q_s(o)$ and such that $\{X_i\}_{0 \leq i \leq n}$ forms a b-bounded anisotropic descending chain. Then $\mathbb{P}\left(A^{(4)}\right) \leq s^2 \left(4b^2\right)^n \lambda^{n+1}/n!$.*

Proof. If we denote by N the number of $n+1$-tuples of pairwise disjoint elements $X_0, \dots, X_n \in$

X satisfying $A^{(4)}$, then using Lemma 4.3 and the Campbell formula we compute

$$\begin{aligned}
\mathbb{E}N &\le \lambda^{n+1} \int \cdots \int 1_{x_0 \in Q_s(o)} 1_{b \ge |x_0 - x_1|_\infty \ge \ldots \ge |x_{n-1} - x_n|_\infty} \mathrm{d}x_n \cdots \mathrm{d}x_0 \\
&= 4\lambda^{n+1} \int \cdots \int 1_{x_0 \in Q_s(o)} 1_{b \ge |x_0 - x_1|_\infty \ge \ldots \ge |x_{n-2} - x_{n-1}|_\infty} \left|x_{n-2} - x_{n-1}\right|_\infty^2 \mathrm{d}x_{n-1} \cdots \mathrm{d}x_0 \\
&= \ldots \\
&= \frac{\left(4b^2\right)^n \lambda^{n+1}}{n!} \int 1_{x_0 \in Q_s(o)} \mathrm{d}x_0 \\
&= \frac{s^2 \left(4b^2\right)^n \lambda^{n+1}}{n!}.
\end{aligned}$$

□

4.2 Proof of Proposition 3.2

In this section we prove Proposition 3.2. We observe that a similar technique as developed in [4] may be applied to obtain a suitable external stabilization result in the current setting. To be more precise, we first prove the following auxiliary result.

Lemma 4.5. *Let $\alpha, s > 0$. Then the probability that there exists a finite set $\xi \subset \mathbb{R}^{2,m} \backslash Q_s^m(o)$ such that $\nu_1 \left(I_{X_o^{(1)} \cup \xi}(x) \cap Q_s(o) \right) \ge s^\alpha$ for some $x \in X_o^{(1)} \cup \xi$ tends to 0 as $s \to \infty$. Here ν_1 denotes the length of a given segment.*

Proof. Without loss of generality we may assume $\alpha < 1$. We consider the cases $x \in \xi$ and $x \in X_o^{(1)}$ separately and start with the case $x \in \xi$. By symmetry it suffices to prove that the probability that there exists a finite $\xi \subset \mathbb{R}^{2,m} \backslash Q_s^m(o)$ such that $\nu_1 \left(I_{X_o^{(1)} \cup \xi}(x) \cap Q_s(o) \right) \ge s^\alpha$ for some $x \in \xi$ with $dir(x) = e_2$ and $\pi_2(x) < 0$ tends to 0 as $s \to \infty$. Put $\delta_1 = s^{-\alpha/2}$, $R_1 = [-\delta_1, \delta_1] \times [-2.5\delta_1, 2.5\delta_1]$ and $R_2 = [0, \delta_1] \times [-\delta_1/2, \delta_1/2]$. For $x \in \mathbb{R}^2$ we denote by $A_x^{(5)}$ the intersection of the events $X^{(1)} \cap ((x + R_2) \times \{-e_1\}) \neq \emptyset$ and $X^{(1)} \cap ((x + R_1) \times M) \setminus ((x + R_2) \times \{-e_1\}) = \emptyset$, see Figure 4 for an illustration.



Figure 4: Occurrence of $A_o^{(5)}$

Then $\mathbb{P}\left(A_x^{(5)}\right) \ge \exp\left(-10\lambda\delta_1^2\right)\left(1 - \exp\left(-\lambda\delta_1^2/4\right)\right) \ge \frac{\delta_1^2}{8}$ for all $x \in \mathbb{R}^2$ and all $s > 0$ sufficiently large. For $\sigma \in \mathbb{R}$ put $M_\sigma^{(3)} = \left\{\sigma e_1 + (-s/2 + 5i\delta_1)e_2 : 1 \le i \le s^\alpha \left(5\delta_1\right)^{-1}\right\}$, so that

$$\mathbb{P}\left(\bigcap_{x \in M_\sigma^{(3)}} \left(A_x^{(5)}\right)^c \right) \le \left(1 - \delta_1^2/8\right)^{s^\alpha (5\delta_1)^{-1}}.$$

Since $s^\alpha\delta_1 = s^{\alpha/2}$, we see that $\mathbb{P}\left(\bigcap_{x\in M^{(3)}_\sigma}\left(A^{(5)}_x\right)^c\right)$ decays sub-exponentially fast as $s\to\infty$. Therefore, also $\mathbb{P}\left(\bigcup_{j=0}^{s\delta^{-1}}\bigcap_{x\in M^{(3)}_{-s/2+j\delta_1}}\left(A^{(5)}_x\right)^c\right)$ decays sub-exponentially fast in s. Note that the occurrence of the event $\bigcup_{x\in M^{(3)}_{-s/2+j\delta_1}} A^{(5)}_x$ implies that for all finite $\xi\subset\mathbb{R}^{2,m}\setminus Q^m_s(o)$ and all $x\in\xi$ with $dir(x)=e_2$, $\pi_2(x)<0$ and $\pi_1(x)\in[-s/2+(j-1)\delta_1, -s/2+j\delta_1]$ we have $\nu_1\left(I_{X^{(1)}_o\cup\xi}(x)\cap Q_s(o)\right)\le s^\alpha$. This proves the first case of the claim.

Next, consider the case $x\in X^{(1)}_o$. Using the Slivnyak-Mecke formula this part can be proven similarly as the case $x\in\xi$, but we include at least some details for the convenience of the reader. Denote by N the number of nodes $x\in X^{(1)}_o$ such that $\nu_1\left(I_{X^{(1)}_o\cup\xi}(x)\cap Q_s(o)\right)\ge s^\alpha$ for some finite $\xi\subset\mathbb{R}^{2,m}\setminus Q^m_s(o)$. Using the Slivnyak-Mecke formula we compute

$$\mathbb{E}\left(N\right)\le\int_{Q_s(o)}\mathbb{P}\left(\exists\xi\subset\mathbb{R}^{2,m}\setminus Q^m_s(o):\nu_1\left(I_{X^{(1)}_o\cup\{x\}\cup\xi}(x)\cap Q_s(o)\right)\ge s^\alpha\right)\mathrm{d}x.$$

By a similar argument as in the previous case we see that the probability

$$\mathbb{P}\left(\exists\xi\subset\mathbb{R}^{2,m}\setminus Q^m_s(o):\nu_1\left(I_{X^{(1)}_o\cup\{x\}\cup\xi}(x)\cap Q_s(o)\right)\ge s^\alpha\right)$$

decays sub-exponentially fast in s. This completes the proof Lemma 4.5. ☐

As immediate corollaries we obtain the absence of trespassing segments and the existence of interior segments.

Corollary 4.6. *Let $s>0$ and write $A^{(6)}_s$ for the event $g_{X^{(1)}}(x)\in Q_s(o)$ for all $x\in\partial^{in}_{X^{(1)}}Q_s(o)$. Then $\lim_{s\to\infty}\mathbb{P}\left(A^{(6)}_s\right)=1$.*

Corollary 4.7. *Let $s>0$ and write $A^{(7)}_s$ for the event $X^{(1)}_o\ne\partial^{out}_{X^{(1)}}Q_s(o)$. Then $\mathbb{P}\left(A^{(7)}_s\right)\to 1$ as $s\to\infty$.*

Furthermore combining Lemma 4.5 with Lemma 4.2 we obtain external stabilization.

Lemma 4.8. *Denote by $A^{(8)}_s$ the event that $\partial^{in}_{X^{(1)}\cap Q^m_{3s}(o)}Q_s(o)=\partial^{in}_{X^{(1)}\cap Q^m_{3s}(o)\cup\xi}Q_s(o)$ for all finite $\xi\subset\mathbb{R}^{2,m}\setminus Q^m_{3s}(o)$. Then $\lim_{s\to\infty}\mathbb{P}\left(A^{(8)}_s\right)=1$.*

Proof. Assume that the event described in Lemma 4.5 is satisfied for $\alpha_1=1/8$ and $3s$ instead of s. Therefore, if the event $A^{(8)}_s$ does not occur, then there exist a finite subset $\xi\subset\mathbb{R}^{2,m}\setminus Q^m_{3s}(o)$ and an element $x\in X^{(1)}\cap Q^m_{2s}(o)$ with $f_{X^{(1)}\cap Q^m_{3s}(o)}(x)\ne f_{X^{(1)}\cap Q^m_{3s}(o)\cup\xi}(x)$. By Lemma 4.2 there exists an $s^{1/4}$-bounded anisotropic descending chain of pairwise disjoint elements of $X^{(1)}\cap Q^m_{3s}(o)\cup\xi$ starting in x and ending in $\mathbb{R}^2\setminus Q_{3s}(o)$. Furthermore, by our assumption this chain consists of at least $s/s^{1/4}=s^{3/4}$ hops. However, by Lemma 4.4 the probability for the occurrence of this event can be bounded from above by $4s^2(4s^{1/2})^{s^{3/4}}/\lfloor s^{3/4}\rfloor!$. Using Stirling's formula, we see that the latter expression tends to 0 as $s\to\infty$. ☐

Finally, consider the property of δ-separatedness.

Lemma 4.9. *Let $s>0$ and put $\delta=s^{-4}$. Write $A^{(9)}_s$ for the event that $|\pi_k(x)-\pi_k(y)|\ge\delta$ for all $x,y\in X^{(1)}\cap Q^m_{3s}(sz)\cup\left\{sz+\frac{s}{2}M\right\}$ with $x\ne y$ and all $k\in\{1,2\}$. Then $\lim_{s\to\infty}\mathbb{P}\left(A^{(9)}_s\right)=1$.*

Proof. For $k\in\{1,2\}$ denote by N_k the number of pairs (x,y) with $x,y\in X^{(1)}\cap Q^m_{3s}(sz)$ and such that $|\pi_k(x)-\pi_k(y)|\le\delta$. Then

$$\begin{aligned}\mathbb{E}N_1&\le\int_{Q_{3s}(o)}\int_{Q_{3s}(o)}1_{|\pi_1(x)-\pi_1(y)|\le\delta}dydx\\&=9s^2\int_{-3s/2}^{3s/2}\int_{-3s/2}^{3s/2}1_{|\eta_1-\xi_1|\le\delta}d\eta_1 d\xi_1\\&=54s^3\delta\\&\le 54s^{-1}.\end{aligned}$$

Similar bounds may be obtained for $\mathbb{E}N_2$ and also for the number of pairs (x, y) with $x \in X^{(1)} \cap Q^m_{3s}(sz)$, $y \in sz + \frac{s}{2}M$ and such that $|\pi_k(x) - \pi_k(y)| \le \delta$. □

Proposition 3.2 now follows by combining Corollaries 4.6 and 4.7 as well as Lemmas 4.8 and 4.9.

4.3 Proofs of Lemmas 3.8 and 3.9

The present section is devoted to the proofs of Lemmas 3.8 and 3.9. We begin by considering a stability result of the lilypond model in a square with respect to changes of those germs whose associated grain does not intersect the square at hand.

Lemma 4.10. *Let* $s > 0$, $z \in \mathbb{Z}^2$ *and* $\varphi^{(1)}, \varphi^{(2)} \in \mathbb{N}'$ *be such that* $\varphi^{(1)}_z \cup \partial^{in}_{\varphi^{(1)}} Q_s(sz) = \varphi^{(2)}_z \cup \partial^{in}_{\varphi^{(2)}} Q_s(sz)$. *Then*

(i) $f_{\varphi^{(1)}}(x) = f_{\varphi^{(2)}}(x)$ *for all* $x \in \varphi^{(1)}_z \cup \partial^{in}_{\varphi^{(1)}} Q_s(sz)$ *with* $\big\{g_{\varphi^{(1)}}(x), g_{\varphi^{(2)}}(x)\big\} \cap Q_s(sz) \neq \emptyset$,

(ii) $g_{\varphi^{(1)}}(x) \in Q_s(sz)$ *for all* $x \in \partial^{in}_{\varphi^{(1)}} Q_s(sz)$ *if and only if* $g_{\varphi^{(2)}}(x) \in Q_s(sz)$ *for all* $x \in \partial^{in}_{\varphi^{(2)}} Q_s(sz)$,

(iii) $\partial^{out}_{\varphi^{(1)}} Q_s(sz) = \partial^{out}_{\varphi^{(2)}} Q_s(sz)$ *and*

(iv) for any finite $\xi \subset Q^m_s(sz)$ *the events* $A^{(2)}_z(\varphi^{(1)}, \xi)$ *and* $A^{(2)}_z(\varphi^{(2)}, \xi)$ *are equivalent.*

Proof. For readability we write f_1, f_2 instead of $f_{\varphi^{(1)}}$, $f_{\varphi^{(2)}}$ (and similarly for $g, I, \partial^{in/out}$). Put $\varphi' = \big\{x \in \varphi^{(1)}_z \cup \partial^{in}_1 Q_s(sz) : \{g_1(x), g_2(x)\} \cap Q_s(sz) \neq \emptyset\big\}$. Our first goal is to show $f_1(x) = f_2(x)$ for all $x \in \varphi'$. Suppose for the sake of deriving a contradiction that there exists $x_1 \in \varphi'$ with $f_1(x_1) \neq f_2(x_1)$, e.g. $f_1(x_1) < f_2(x_1)$. By Lemma 4.2 it suffices to show that for any such $x_1 \in \varphi'$ we have $x_2 \in \varphi'$, where $x_2 = h_1(x_1)$.

First observe that $g_1(x_1) \in Q_s(sz)$. Indeed, the contrary would imply $g_2(x_1) \in Q_s(sz)$ and $x_1 \in \partial^{in}_2 Q_s(sz)$, but $x_1 \notin \partial^{in}_1 Q_s(sz)$ contradicting our assumption $\partial^{in}_1 Q_s(sz) = \partial^{in}_2 Q_s(sz)$. In particular, this shows $x_2 \in \varphi^{(1)}_z \cup \partial^{in}_1 Q_s(sz)$. From the assumption $f_1(x_1) < f_2(x_1)$ we then conclude $f_2(x_2) < f_1(x_2)$. We claim that $g_2(x_2) \in Q_s(sz)$ and assume the contrary for the sake of deriving a contradiction. Then we conclude from $g_2(x_2) \in [x_2, g_1(x_1)]$ and $g_1(x_2) \in [g_1(x_1), g_2(x_2)]$ that $x_2 \notin Q_s(sz)$, $x_2 \notin \partial^{in}_2 Q_s(sz)$ and $x_2 \in \partial^{in}_1 Q_s(sz)$, contradicting our assumption. This completes the proof of $f_1(x) = f_2(x)$ for all $x \in \varphi'$. Assertion (ii) is an immediate consequence of assertion (i).

To prove the third claim, let $x \in \varphi^{(1)}_z$. If $x \notin \partial^{out}_1 Q_s(sz)$, then $g_1(x) \in Q_s(sz)$ and therefore also $g_2(x) = g_1(x) \in Q_s(sz)$. In other words $\varphi^{(1)}_z \setminus \partial^{out}_1 Q_s(sz) \subset \varphi^{(1)}_z \setminus \partial^{out}_2 Q_s(sz)$ and the other inclusion follows by symmetry. Finally, the last claim is an immediate consequence of assertion (iii), since the event $A^{(2)}_z(\varphi^{(1)}, \xi)$ depends on $\varphi^{(1)}$ only through $\varphi^{(1)}_z$, $\partial^{in}_1 Q_s(sz)$ and $\partial^{out}_1 Q_s(sz)$. □

Using Lemma 4.10 we can now provide a proof for Lemma 3.8

Proof of Lemma 3.8. Item (i) is an immediate consequence of the external stabilization property of s-good sites. In particular, we may apply Lemma 4.10 to $\varphi^{(1)} = X^{(1)}$ and $\varphi^{(2)} = X^{(3)}$ to obtain $f_{X^{(3)}}(x) = f_{X^{(1)}}(x)$ for all $x \in X^{(3)}$ with $\{g_{X^{(3)}}(x), g_{X^{(1)}}(x)\} \cap Q_s(sz) \neq \emptyset$. □

In the next two lemmas we consider the precise effects of adding nice sprinkling inside a given square.

Lemma 4.11. *Let* $s, \delta > 0$, $z \in \mathbb{Z}^2$, $\varphi \in \mathbb{N}'$, $\xi \subset Q_s(sz)$ *finite and put* $\delta = s^{-4}$. *Furthermore, assume that* $\varphi_z \cup \partial^{in}_\varphi Q_s(sz) \cup \{sz + \frac{s}{2}M\}$ *is* δ*-separated and that* $A^{(2)}_z(\varphi, \xi)$ *occurs. Then the following items are satisfied.*

(i) $f_{\varphi \cup \xi}(x) \le \frac{3\delta}{8}$ *for all* $x \in \varphi_z \setminus \partial^{out}_\varphi Q_s(sz)$

(ii) $[x, g_{\varphi \cup \xi}(x) - \frac{3\delta}{8} dir(x)] \cap Q_s(sz) = \emptyset$ *for all* $x \in \partial^{in}_\varphi Q_s(sz)$

Proof. Let $x \in \varphi_z \setminus \partial^{out}_{\varphi} Q_s(sz)$ and let $D \subset \xi$ be a $(y, \delta/16)$-cycle, where $y = x + \frac{3\delta}{8} dir(x)$. Observe that by δ-separatedness and Lemma 4.11 we have $f_{\varphi \cup \xi}(x') = f_D(x')$ for all $x' \in D$, so that $f_{\varphi \cup \xi}(x) > \frac{3\delta}{8}$ would contradict the hard-core property. Similarly, let $x \in \partial^{in}_{\varphi} Q_s(sz)$ and let $D \subset \xi$ be a $(y, \delta/16)$-cycle, where $y = P_x + \frac{3\delta}{8} dir(x)$ and P_x denotes the first intersection point of $I_{\varphi}(x)$ and $Q_s(sz)$. Observe that by δ-separatedness and Lemma 4.11 we have $f_{\varphi \cup \xi}(x') = f_D(x')$ for all $x' \in D$, so that again $g_{\varphi \cup \xi}(x) - \frac{3\delta}{8} dir(x) \in Q_s(sz)$ would contradict the hard-core property. □

Lemma 4.12. *Let $s > 0$, $z \in \mathbb{Z}^2$, let $\xi \subset Q^m_s(sz)$ be finite and let $\varphi \in \mathbb{N}'$. Assume*

- *(i)* $\varphi_z \cup \partial^{in}_{\varphi} Q_s(sz) \cup \{sz + \frac{s}{2} M\}$ *is δ-separated,*
- *(ii)* $g_{\varphi}(x) \in Q_s(sz)$ *for all* $x \in \partial^{in}_{\varphi} Q_s(sz)$ *and*
- *(iii)* $A^{(2)}_z(\varphi, \xi)$ *occurs.*

Then

- *(i)* $f_{\varphi \cup \xi}(x) = f_{\xi}(x)$ *for all* $x \in \xi$,
- *(ii)* $f_{\varphi \cup \xi}(x) = f_{\varphi}(x)$ *for all* $x \in \varphi$ *with* $\{g_{\varphi}(x), g_{\varphi \cup \xi}(x)\} \not\subset Q_s(sz)$,
- *(iii)* $\partial^{in}_{\varphi} Q_s(sz') = \partial^{in}_{\varphi \cup \xi} Q_s(sz')$ *for all* $z' \in \mathbb{Z}^2$ *and*
- *(iv)* $h_{\varphi \cup \xi}(x) \in \xi$ *for all* $x \in \varphi$ *with* $g_{\varphi}(x) \in Q_s(sz)$.

Proof. Assertion (i) follows immediately from δ-separatedness and Lemma 3.4. Next we claim that if $x \in \varphi$ is such that $g_{\varphi}(x) \in Q_s(sz)$ then $f_{\varphi \cup \xi}(x) \le f_{\varphi}(x)$ and if additionally $I_{\varphi \cup \xi}(x) \cap Q_s(sz) \ne \emptyset$, then $h_{\varphi \cup \xi}(x) \in \xi$. To prove this claim we distinguish two cases. If $x \notin Q_s(sz)$, then $x \in \partial^{in}_{\varphi} Q_s(sz)$. Therefore, Lemma 4.11 shows $[x, g_{\varphi \cup \xi}(x_1) - 3\delta/8 \cdot dir(x)] \cap Q_s(sz) = \emptyset$, so that $f_{\varphi \cup \xi}(x) - 3\delta/8 \le f_{\varphi}(x)$. Hence, $f_{\varphi}(x) \le f_{\varphi \cup \xi}(x)$ would yield a contradiction to δ-separatedness of $\varphi_z \cup \partial^{in}_{\varphi} Q_s(sz) \cup \{sz + \frac{s}{2} M\}$. Additionally, if $I_{\varphi \cup \xi}(x) \cap Q_s(sz) \ne \emptyset$ then δ-separatedness can also be used to deduce $h_{\varphi \cup \xi}(x) \in \xi$. Similarly, if $x \in Q_s(sz)$, then $x \in \varphi_z \setminus \partial^{out}_{\varphi} Q_s(sz)$, so that from $A^{(2)}_z(\varphi, \xi)$ and Lemma 4.11 we conclude $f_{\varphi \cup \xi}(x) \le 3\delta/8$. In particular, δ-separatedness of $\varphi_z \cup \partial^{in}_{\varphi} Q_s(sz)$ again implies $f_{\varphi}(x) \ge f_{\varphi \cup \xi}(x)$ and $h_{\varphi \cup \xi}(x) \in \xi$.

To prove item (ii) put $\varphi' = \{x \in \varphi : \{g_{\varphi}(x), g_{\varphi \cup \xi}(x)\} \not\subset Q_s(sz)\}$. Then we show $f_{\varphi}(x) = f_{\varphi \cup \xi}(x)$ for all $x \in \varphi'$. Suppose for the sake of deriving contradiction that there exists $x_1 \in \varphi'$ with $f_{\varphi}(x_1) \ne f_{\varphi \cup \xi}(x_1)$. By Lemma 4.2 it suffices to show that for any such $x_1 \in \varphi'$ we have $h_{\varphi}(x_1) \in \varphi'$ if $f_{\varphi}(x_1) < f_{\varphi \cup \xi}(x_1)$ and $h_{\varphi \cup \xi}(x_1) \in \varphi'$ if $f_{\varphi \cup \xi}(x_1) < f_{\varphi}(x_1)$.

case $f_{\varphi}(x_1) < f_{\varphi \cup \xi}(x_1)$**.** Put $x_2 = h_{\varphi}(x_1)$. If $g_{\varphi}(x_1) \notin Q_s(sz)$ then we deduce $x_2 \in \varphi'$ from $g_{\varphi}(x_1) \in [g_{\varphi \cup \xi}(x_2), g_{\varphi}(x_2))$. But if $g_{\varphi}(x_1) \in Q_s(sz)$ then $f_{\varphi}(x_1) < f_{\varphi \cup \xi}(x_1)$ yields a contradiction to the argument presented at the beginning of the proof.

case $f_{\varphi \cup \xi}(x_1) < f_{\varphi}(x_1)$**.** Put $x_2 = h_{\varphi \cup \xi}(x_1)$. Again if $g_{\varphi \cup \xi}(x_1) \notin Q_s(sz)$ then we deduce $x_2 \notin \xi$ and deduce further from $g_{\varphi \cup \xi}(x_1) \in [g_{\varphi}(x_2), g_{\varphi \cup \xi}(x_2))$ that $x_2 \in \varphi'$. Therefore, we may restrict our attention to the case $g_{\varphi \cup \xi}(x_1) \in Q_s(sz)$, so that $g_{\varphi}(x_1) \notin Q_s(sz)$. From assumption (ii) we conclude $x_1 \in Q_s(sz)$ which implies $x_1 \in \partial^{out}_{\varphi} Q_s(sz)$. By assumption (iii) this implies $x_2 \in \varphi$, so that $f_{\varphi}(x_2) < f_{\varphi \cup \xi}(x_2)$. This shows $x_2 \in \varphi'$, since the contrary would yield a contradiction to the argument presented at the beginning of the proof.

To show item (iii) assume the contrary and then choose $z' \in \mathbb{Z}^2$ and an element x in the symmetric difference $\partial^{in}_{\varphi} Q_s(sz') \Delta \partial^{in}_{\varphi \cup \xi} Q_s(sz')$. By assertion (i) this implies $x \in \varphi$ and by item (ii) we conclude that both $g_{\varphi}(x) \in Q_s(sz)$ and $g_{\varphi \cup \xi}(x) \in Q_s(sz)$. But then, trivially, $x \in \partial^{in}_{\varphi} Q_s(sz')$ iff $x \in \partial^{in}_{\varphi \cup \xi} Q_s(sz')$.

Finally, to prove item (iv) let $x \in \varphi$ with $g_{\varphi}(x) \in Q_s(sz)$ be arbitrary. In particular, by item (iii) of the assertion we have $I_{\varphi \cup \xi}(x) \cap Q_s(sz) \ne \emptyset$. The argument at the beginning of the proof therefore implies $h_{\varphi \cup \xi}(x) \in \xi$. □

Using these deterministic auxiliary results, we can now provide the proof of Lemma 3.9.

Proof of Lemma 3.9. Naturally, we prove the claim by induction on $j \geq 0$. When $j = 0$ items $(ii), (iii), (v)$ are trivial, while items (i) and (iv) follow from Lemma 3.8.

Next suppose that claims (i) to (v) have already been proven for $j \geq 0$; we explain how to deduce the result for $j + 1$. Observe that all claims follow trivially from induction hypothesis when $Y_{z_{j+1}} = 0$, so without loss of generality we may assume $Y_{z_{j+1}} = 1$. First note that by induction hypothesis $\partial^{in}_{X^{(3,j)}} Q_s(sz) = \partial^{in}_{X^{(1)}} Q_s(sz)$ for all $z \in R^c$, so that we may apply Lemma 4.10 with $z = z_{j+1}$, $\varphi^{(1)} = X^{(1)}$ and $\varphi^{(2)} = X^{(3,j)}$. Hence, we obtain $A^{(2)}_{z_{j+1}}\left(X^{(3,j)}, X^{(2)}_{z_{j+1}}\right)$ and $g_{X^{(3,j)}}(x) \in Q_s(sz_{j+1})$ for all $x \in \partial^{in}_{X^{(3,j)}} Q_s(sz_{j+1}) = \partial^{in}_{X^{(1)}} Q_s(sz_{j+1})$. From Lemma 4.12 we therefore conclude

(i) $f_{X^{(3,j+1)}}(x) = f_{X^{(2)}_{z_{j+1}}}(x)$ for all $x \in X^{(2)}_{z_{j+1}}$,

(ii) $f_{X^{(3,j+1)}}(x) = f_{X^{(3,j)}}(x)$ for all $x \in X^{(3,j)}$ with $g_{X^{(3,j)}}(x) \notin Q_s(sz_{j+1})$ and

(iii) $\partial^{in}_{X^{(3,j+1)}} Q_s(sz') = \partial^{in}_{X^{(3,j)}} Q_s(sz')$ for all $z' \in \mathbb{Z}^2$.

(iv) $h_{X^{(3,j+1)}}(x) \in X^{(2)}_{z_{j+1}}$ for all $x \in X^{(3,j)}$ with $g_{X^{(3,j)}}(x) \in Q_s(sz_{j+1})$

Item (iii) above in conjunction with induction hypothesis then proves item (i) of the original assertion. Furthermore, for $z \in R$ and all $x \in X^{(3)}$ with $g_{X^{(3)}}(x) \in Q_s(sz)$ the induction hypothesis implies $g_{X^{(3,j)}}(x) \in Q_s(sz)$ so that item (ii) above may be used to deduce item (ii) of the original claim. Item (i) above also shows item (iii) of the original claim in the special case $z = z_{j+1}$. If $z \neq z_{j+1}$ and $z \in R^c$, then we conclude from induction hypothesis and item (ii) above that for $x \in X^{(2,j+1)}_z = X^{(2,j)}_z$ we have

$$\begin{aligned} f_{X^{(3,j+1)}}(x) &= f_{X^{(3,j)}}(x) \\ &= f_{X^{(2,j)}_z}(x). \end{aligned}$$

This completes the proof of item (iii) and it remains to prove items (iv) and (v). We consider again first the case $z = z_{j+1}$ (where $X^{(2)}_{z_{j+1}} \neq \emptyset$). By item (iv) above it suffices to show that $g_{X^{(3)}}(x) \in Q_s(sz_{j+1})$ implies $g_{X^{(3,j)}}(x) \in Q_s(sz_{j+1})$. Observe that by induction $g_{X^{(1)}}(x) = g_{X^{(3,j)}}(x)$, while Lemma 3.8 implies that $g_{X^{(1)}}(x) = g_{X^{(3)}}(x)$, thereby completing the proof of item (iv) in the case $z = z_{j+1}$. Next, let $z \neq z_{j+1}$, $z \in R^c$ and $g_{X^{(3)}}(x) \in Q_s(sz)$. First we consider item (iv) and hence assume $X^{(2,j)}_z = \emptyset$. The induction hypothesis yields $g_{X^{(3,j)}}(x) = g_{X^{(1)}}(x) \in Q_s(sz)$, so that item (ii) above and shows $f_{X^{(3,j+1)}}(x) = f_{X^{(3,j)}}(x) = f_{X^{(1)}}(x)$. Finally, assume $X^{(2,j)}_z \neq \emptyset$. Here the induction hypothesis yields $h_{X^{(3,j)}}(x) \in X^{(2)}_z \subset Q_s(sz)$, so that $g_{X^{(3,j)}}(x) \in Q_s(sz)$. By item (ii) above we therefore obtain $h_{X^{(3,j+1)}}(x) = h_{X^{(3,j)}}(x) \in X^{(2)}_z$. ☐

Acknowledgments

The author is grateful to D. Daley for proposing the percolation problem. The author also thanks D. Daley, V. Schmidt and S. Ziesche for interesting discussions as well as useful remarks on an earlier version of this manuscript. This work has been supported by a research grant from DFG Research Training Group 1100 at Ulm University.

References

[1] M. Aizenman, J. Chayes, L. Chayes, J. Fröhlich, and L. Russo. On a sharp transition from area law to perimeter law in a system of random surfaces. *Communications in Mathematical Physics*, 92:19–69, 1983.

[2] D. J. Daley and G. Last. Descending chains, the lilypond model, and mutual-nearest-neighbour matching. *Advances in Applied Probability*, 37:604–628, 2005.

[3] D. J. Daley, H. Stoyan, and D. Stoyan. The volume fraction of a Poisson germ model with maximally non-overlapping spherical grains. *Advances in Applied Probability*, 31:610–624, 1999.

[4] S. Ebert. *Stochastische Modellierung einer Klasse wachstumsmaximaler Keim-Korn-Modelle*. PhD thesis, KIT, February 2012.

[5] O. Häggström and R. Meester. Nearest neighbor and hard sphere models in continuum percolation. *Random Structures Algorithms*, 9:295–315, 1996.

[6] M. Heveling and G. Last. Existence, uniqueness, and algorithmic computation of general lilypond systems. *Random Structures Algorithms*, 29:338–350, 2006.

[7] G. Last and M. D. Penrose. Percolation and limit theory for the Poisson lilypond model. *Random Structures Algorithms*, 42:226–249, 2013.

[8] T. M. Liggett, R. H. Schonmann, and A. M. Stacey. Domination by product measures. *Annals of Probability*, 25:71–95, 1997.